



Acer campestre



Magasság	circa 12 m
Szélesség	7-8m
Korona	oval, dark, dense crown, capricious growing
Kéreg és ágak	grey bark with conspicuous corky fissures
Leaf	3/5 lobes, dark green, 6 - 10 (12)cm
Őszi színek	sárga
Virágok	small, terminal corymbs, yellow-green, May
Gyümölcsök	single-seed winged nutlets, always paired
Tüskés/tövises	egyik sem
Mérgező tulajdonság	rendszerint nem mérgező emberek, (nagytestű) háziállatok és haszonállatok számára
Talajtípus	any soil, except dry infertile sandy soil
Talajnedvesség	rövid idejű árvíznek ellenáll
Ültetési hely követelményei	Elviseli a laza útburkolatot
télállóság	5a (-28,8 és -26,1 °C)
Szélellenállóság	very good
Egyéb tűrőképességek	fagytűrő (télállósági fok: 1-6), széltűrő, útsónak ellenáll
Fauna fák	fagytűrő (télállósági fok: 1-6), széltűrő, útsónak ellenáll, gazdanövény pillangók számára
Alkalmazások	sugárutak és széles utcák, faültető edények, tetőkeretek, tengerpart, ipari terület
Forma	magas törzsű fa, tőtől elágazó fák, többtörzsű fa
Eredet	Europe, western Asia and North Africa

A medium sized tree up to 12 m high. The crown is a-symmetric because of the irregular growth of the branches. The branches have conspicuous corky fissures, both on mature and young wood. Juvenile foliage may have a reddish look when it unfurls, but soon turns dark green. The colour in autumn is dark yellow and last for a long time. In the Netherlands *Acer campestre* is indigenous and has been used by man for a very long time. Not only as a tree, but also pruned as a hedge this plant is very functional. Because of its shade-tolerance and ability to grow on industrial estates it is a very multi-functional plant. Forms a strongly branching main root with very many rootlets. The plant is propagated by seeding and therefore variations in the off-spring can occur. This is visible because some features (autumn colour, corky fissures on the branches) are present to a more or lesser extent. *Acer campestre* is often used in hedges since it stands up exceptionally well to pruning. But pruning should never be done in the spring because the tree's sap rises early.